



Hexaaquazinc(II) dinitrate bis[5-(pyridinium-3-yl)tetrazol-1-ide]

Ignacio Chi-Duran, Javier Enriquez, Andres Vega, Felipe Herrera and
Dinesh Pratap Singh

Acta Cryst. (2018). E74, 1231–1234



IUCr Journals

CRYSTALLOGRAPHY JOURNALS ONLINE

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.





Hexaaquazinc(II) dinitrate bis[5-(pyridinium-3-yl)tetrazol-1-ide]

Ignacio Chi-Duran,^a Javier Enriquez,^a Andres Vega,^b Felipe Herrera^{a,c} and Dinesh Pratap Singh^{a,c,*}

^aDepartment of Physics, University of Santiago, Av. Ecuador 3493, Estación Central, Santiago, Chile, ^bDepartamento de Ciencias Químicas, Universidad Nacional Andres Bello, Av Republica 275 3er Piso, Santiago, Region Metropolitana, Chile, and ^cMillennium Institute for Research in Optics (MIRO), Chile. *Correspondence e-mail: singh.dinesh@usach.cl

Received 29 May 2018

Accepted 3 August 2018

Edited by A. M. Chippindale, University of Reading, England

Keywords: crystal structure; pyridin-3-yltetrazole; hexaaquazinc(II) complex; π - π stacking; hydrogen-bonding.



CCDC reference: 1860162

Supporting information: this article has supporting information at journals.iucr.org/e

Hexaaquazinc(II) dinitrate 5-(pyridinium-3-yl)tetrazol-1-ide, $[\text{Zn}(\text{H}_2\text{O})_6](\text{NO}_3)_2 \cdot 2\text{C}_6\text{H}_5\text{N}_5$, crystallizes in the space group $P\bar{1}$. The asymmetric unit contains one zwitterionic 5-(pyridinium-3-yl)tetrazol-1-ide molecule, one NO_3^- anion and one half of a $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$ cation ($\bar{1}$ symmetry). The pyridinium and tetrazolide rings in the zwitterion are nearly coplanar, with a dihedral angle of $5.4(2)^\circ$. Several $\text{O}-\text{H} \cdots \text{N}$ and $\text{N}-\text{H} \cdots \text{O}$ hydrogen-bonding interactions exist between the $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$ cation and the N atoms of the tetrazolide ring, and between the nitrate anions and the N-H groups of the pyridinium ring, respectively, giving rise to a three-dimensional network. The 5-(pyridinium-3-yl)tetrazol-1-ide molecules show parallel-displaced π - π stacking interactions; the centroid-centroid distance between adjacent tetrazolide rings is $3.6298(6) \text{ \AA}$ and that between the pyridinium and tetrazolide rings is $3.6120(5) \text{ \AA}$.

1. Chemical context

Tetrazole functional groups have attracted increased attention in recent years due to their use in drug design and their employment as isosteric substituents of carboxylic acids (Herr, 2002), as well as their ability to produce a large variety of metal-organic frameworks (MOFs) (Zhao *et al.*, 2008; Chi-Duran *et al.*, 2018). Push-pull tetrazole complexes with both electron-donor and electron-acceptor substituents have shown efficient second-order nonlinear optical activity in powdered samples (Masahiko *et al.*, 1994), ferroelectric behaviour (Liu *et al.*, 2015) and strong photoluminescence (Zhang *et al.*, 2014). The *in-situ* synthesis of tetrazole compounds can be realized by the Demko-Sharpless method, in which zinc salts catalyze the cycloaddition reaction between sodium azide and nitrile compounds to form the tetrazole ring (Demko & Sharpless, 2001). In this work, pyridyltetrazole, synthesized at low pH using the Demko-Sharpless method, is cocrystallized in the presence of $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$ and NO_3^- ions, to obtain the title compound (Fig. 1).



2. Structural commentary

The asymmetric unit of the title compound is composed of one 5-(pyridinium-3-yl)tetrazol-1-ide zwitterion, one NO_3^- anion and one half of a $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$ cation. The hexaaquazinc(II)

Table 1
Selected geometric parameters (Å, °).

Zn1—O3	2.0353 (11)	Zn1—O2 ⁱ	2.1011 (12)
Zn1—O3 ⁱ	2.0354 (11)	Zn1—O1 ⁱ	2.1841 (11)
Zn1—O2	2.1011 (12)	Zn1—O1	2.1841 (11)
O3—Zn1—O2	90.01 (5)	O2—Zn1—O1 ⁱ	92.10 (5)
O3—Zn1—O2 ⁱ	89.99 (5)	O3—Zn1—O1	89.47 (4)
O3—Zn1—O1 ⁱ	90.53 (4)	O2—Zn1—O1	87.90 (5)

Symmetry code: (i) $-x, -y + 2, -z + 2$.

complex exhibits regular octahedral geometry (Table 1), and the tetrazolide and pyridinium rings of the zwitterion are close to being coplanar, with a dihedral angle of $5.4(2)^\circ$ (Fig. 2). The geometric parameters of the tetrazolide ring are comparable to those in other reported tetrazole compounds (Mu *et al.*, 2010; Dai & Chen, 2011*a,b*). The H atom attached to the N atom of the pyridine ring could not be located in the Fourier density map. Therefore, the H atom was placed in accordance with similar reported structures containing $[\text{Mg}(\text{H}_2\text{O})_6]\text{X}_2$ ($\text{X} = \text{Cl}^-, \text{Br}^-$) cocrystallized with 5-(pyridinium-3-yl)tetrazolide (Dai & Chen, 2011*a,b*).

3. Supramolecular features

A three-dimensional network of hydrogen bonds involving the pyridinium–tetrazolide zwitterions, hexaaquazinc(II) complex cations and nitrate ions serves to hold the structure together (Table 2 and Fig. 3). The N atoms of the tetrazole ring interact




Figure 1
The molecular structure of the asymmetric unit (plus the three water molecules of the hexaaquazinc cation generated by symmetry), showing the atom labelling and displacement ellipsoids drawn at the 50% probability level. [Symmetry code: (i) $-x, 2 - y, 2 - z$.]

Table 2
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1w ⁱⁱ ⋯O5 ⁱⁱ	0.85	1.96	2.8067 (17)	172
O1—H2w ⁱⁱⁱ ⋯N1 ⁱⁱⁱ	0.85	1.96	2.8029 (17)	173
O2—H3w ^{iv} ⋯N4 ^{iv}	0.85	2.02	2.8589 (17)	170
O2—H4w ^v ⋯O1 ^v	0.85	2.08	2.9228 (17)	171
O3—H5w ^{vi} ⋯N3 ^{vi}	0.85	1.91	2.7446 (17)	168
O3—H6w ^{vi} ⋯N1 ^{vi}	0.85	2.72	3.4294 (17)	142
O3—H6w ^{vii} ⋯N2 ^{vii}	0.85	1.97	2.8076 (17)	169
N5—H5N ^{vii} ⋯N6 ^{vii}	0.82	2.61	3.344 (2)	149
N5—H5N ^{viii} ⋯O4 ^{viii}	0.82	1.92	2.7384 (18)	173
N5—H5N ^{viii} ⋯O5 ^{viii}	0.82	2.62	3.1347 (19)	123
C4—H4 ^{viii} ⋯O5 ^{viii}	0.93	2.65	3.452 (2)	145
C5—H5 ^{ix} ⋯O4 ^{ix}	0.93	2.52	3.292 (2)	141
C5—H5 ^{ix} ⋯O6 ^{ix}	0.93	2.52	3.422 (3)	165
C6—H6 ^{vii} ⋯O5 ^{vii}	0.93	2.41	3.047 (2)	126

Symmetry codes: (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x + 1, -y + 2, -z + 1$; (iv) $x, y + 1, z + 1$; (v) $-x + 1, -y + 2, -z + 2$; (vi) $x - 1, y, z + 1$; (vii) $x - 1, y, z$; (viii) $x - 1, y + 1, z$; (ix) $-x, -y + 1, -z + 1$.

with the octahedral complex, $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$, through $\text{O}\cdots\text{H}\cdots\text{N}$ hydrogen bonds, exhibiting $D\cdots A$ distances in the range 2.7446 (17)–2.8589 (17) Å. Additionally, the pyridinium ring is involved in $\text{N}\cdots\text{H}\cdots\text{O}$ hydrogen bonding to nitrate atom O4, with an $\text{N}\cdots\text{O}$ distance of 2.7384 (18) Å. These interactions are shown in the crystal packing diagram (Fig. 3). The structure also shows parallel-displaced π – π stacking interactions, which arise from partial overlap between the




Figure 2
Partial crystal packing of the title compound, showing the hydrogen-bonding interactions between $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$ and the tetrazolide ring. [Symmetry codes: (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x + 1, -y + 2, -z + 1$; (v) $x, y - 1, z - 1$; (xi) $x + 1, y, z - 1$.]

Table 3
 π - π stacking interaction lengths (Å).

Cg1 and Cg2 are the centroids of the C1/N1/N2/N3/N4 and C2-C6/N5 rings, respectively.

Centroid-centroid	Distance	Tetrazolide interplane distance
Cg1-Cg1 ⁱⁱ	3.6298 (6)	3.23 (1)
Cg1-Cg2 ^{vii}	3.6120 (5)	3.10 (3)

Symmetry codes: (ii) $-x + 1, -y + 1, -1 + z$; (vii) $x - 1, y, z$.

tetrazolide and pyridinium rings in adjacent zwitterions, and extend along the a axis parallel to the (010) plane. These parallel-displaced π - π interactions lead to interplanar distances of 3.21 (1) and 3.10 (3) Å, and two centroid-centroid distances (Table 3). The centroid-centroid distance between the tetrazolide groups is 3.6298 (6) Å and between the pyridinium and tetrazolide rings is 3.6120 (5) Å (Table 3 and Fig. 4).

4. Database survey

We found two previously reported structures that are closely related to the title compound. They both involve a hexa-aquamagnesium(II) cation with a halide counter-ion [chloride (Dai & Chen, 2011*b*) or bromide (Dai & Chen, 2011*a*)] cocrystallized in the presence of 5-(pyridinium-3-yl)tetrazol-1-ide zwitterions (Dai & Chen, 2011*a,b*). There are more hydrogen-bonding interactions in our compound than in the [Mg(H₂O)₆]X₂·2C₆H₅N₅ structures, as more hexa-aquazinc(II) complexes can interact with the N atoms of the tetrazole units. Parallel-displaced π - π stacking interactions occur in the title compound and in [Mg(H₂O)₆]X₂·2C₆H₅N₅. In [Mg(H₂O)₆]Cl₂·2C₆H₅N₅, the pyridinium-tetrazolide zwitterions have alternating orientations in the supramolecular arrangement, whereas in the title compound, the zwitterions are oriented in the same direction, allowing a possible coupling transition




Figure 3
 The crystal packing of the title compound, viewed along the [100] direction, showing O-H...N and N-H...O interactions (cyan lines).




Figure 4
 Partial crystal packing, showing π - π interactions between tetrazole and pyridinium rings, with $d_1 = 3.6298$ (6) Å and $d_2 = 3.6120$ (5) Å. [Symmetry codes: (ii) $-x + 1, -y + 1, -1 + z$; (vii) $x - 1, y, z$.]

between dipole moments similar to J-aggregates (Spano, 2010).

Table 4
 Experimental details.

Crystal data	
Chemical formula	[Zn(H ₂ O) ₆](NO ₃) ₂ ·2C ₆ H ₅ N ₅
M_r	591.81
Crystal system, space group	Triclinic, $P\bar{1}$
Temperature (K)	293
a, b, c (Å)	5.6582 (11), 8.4632 (16), 12.046 (2)
α, β, γ (°)	97.209 (2), 91.123 (2), 93.949 (2)
V (Å ³)	570.67 (19)
Z	1
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	1.16
Crystal size (mm)	0.49 × 0.21 × 0.09
Data collection	
Diffractometer	Bruker SMART CCD area detector
Absorption correction	Numerical (SADABS; Bruker, 2008)
T_{\min}, T_{\max}	0.742, 0.903
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	4429, 2217, 2132
R_{int}	0.013
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.617
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.022, 0.054, 1.08
No. of reflections	2217
No. of parameters	198
No. of restraints	13
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³)	0.37, -0.35

Computer programs: SMART and SAINT (Bruker, 2008), SHELXTL (Sheldrick, 2008) and SHELXL2014 (Sheldrick, 2015).

5. Synthesis and crystallization

All the reactants and chemicals were purchased from Sigma Aldrich and utilized without further purification. A mixture of 3-cyanopyridine (4 mmol), NaN_3 (6 mmol) and ZnCl_2 (2 mmol) were dissolved in 6 ml of distilled water. This mixture was transferred to a glass bottle and then heated at 378 K for 24 h. The pH was adjusted using a HNO_3 (66%) solution immediately after mixing the reactants, and was monitored with a pH meter (pH2700 Oakton) until reaching a pH of 2.0. The reaction mixture was then cooled to 318 K and kept at this temperature for 16 h. The colourless block-shaped crystals obtained were washed with ethanol to give 353 mg (yield 30%) of the title compound.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 4. All H atoms bonded to C atoms were positioned geometrically and treated as riding atoms, using $\text{C}-\text{H} = 0.93 \text{ \AA}$ and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Moreover, all H atoms in the hexaaquazinc(II) complex were refined with a distance restraint of $\text{O}-\text{H} = 0.85 \text{ \AA}$ and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

Funding information

Funding for this research was provided by: Fondecyt Regular (award No. 1151527); Proyecto REDES ETAPA INICIAL,

Convocatoria 2017 (award No. REDI170423); Millennium Institute for Research in Optics (MIRO); Basal USA (award No. 1799).

References

- Bruker (2008). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chi-Duran, I., Enriquez, J., Manquian, C., Wrighton-Araneda, K., Cañon-Mancisor, W., Venegas-Yazigi, D., Herrera, F. & Pratap Singh, D. (2018). *ACS Omega*, **3**, 801–807.
- Dai, J. & Chen, X.-Y. (2011a). *Acta Cryst.* **E67**, m113.
- Dai, J. & Chen, X.-Y. (2011b). *Acta Cryst.* **E67**, m171.
- Demko, Z. P. & Sharpless, K. B. (2001). *J. Org. Chem.* **66**, 7945–7950.
- Herr, R. J. (2002). *Bioorg. Med. Chem.* **10**, 3379–3393.
- Liu, D.-S., Sui, Y., Chen, W.-T. & Feng, P. (2015). *Cryst. Growth Des.* **15**, 4020–4025.
- Masahiko, S., Hideki, I., Shinichi, Y., Fumihide, F., Masanao, E., Katsuya, W. & Nobuo, S. (1994). *Jpn J. Appl. Phys.* **33**, 169–170.
- Mu, Y.-Q., Zhao, J. & Li, C. (2010). *Acta Cryst.* **E66**, m1667.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Spano, F. C. (2010). *Acc. Chem. Res.* **43**, 429–439.
- Zhang, Q., Chen, D., He, X., Huang, S., Huang, J., Zhou, X., Yang, Z., Li, J., Li, H. & Nie, F. (2014). *CrystEngComm*, **16**, 10485–10491.
- Zhao, H., Qu, Z.-R., Ye, H.-Y. & Xiong, R.-G. (2008). *Chem. Soc. Rev.* **37**, 84–100.

supporting information

Acta Cryst. (2018). E74, 1231-1234 [https://doi.org/10.1107/S205698901801112X]

Hexaaquazinc(II) dinitrate bis[5-(pyridinium-3-yl)tetrazol-1-ide]

Ignacio Chi-Duran, Javier Enriquez, Andres Vega, Felipe Herrera and Dinesh Pratap Singh

Computing details

Data collection: *SMART* (Bruker, 2008); cell refinement: *SMART* (Bruker, 2008); data reduction: *SAINT* (Bruker, 2008); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Hexaaquazinc(II) dinitrate bis[5-(pyridinium-3-yl)tetrazol-1-ide]

Crystal data

$[\text{Zn}(\text{H}_2\text{O})_6](\text{NO}_3)_2 \cdot 2\text{C}_6\text{H}_5\text{N}_5$

$M_r = 591.81$

Triclinic, $P\bar{1}$

$a = 5.6582$ (11) Å

$b = 8.4632$ (16) Å

$c = 12.046$ (2) Å

$\alpha = 97.209$ (2)°

$\beta = 91.123$ (2)°

$\gamma = 93.949$ (2)°

$V = 570.67$ (19) Å³

$Z = 1$

$F(000) = 304$

$D_x = 1.722$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7831 reflections

$\theta = 2.8\text{--}29.5^\circ$

$\mu = 1.16$ mm⁻¹

$T = 293$ K

Block, colorless

$0.49 \times 0.21 \times 0.09$ mm

Data collection

Bruker SMART CCD area detector
diffractometer

Radiation source: sealed tube

phi and ω scans

Absorption correction: numerical
(SADABS; Bruker, 2008)

$T_{\min} = 0.742$, $T_{\max} = 0.903$

4429 measured reflections

2217 independent reflections

2132 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.013$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 2.4^\circ$

$h = -6 \rightarrow 6$

$k = -10 \rightarrow 10$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.022$

$wR(F^2) = 0.054$

$S = 1.08$

2217 reflections

198 parameters

13 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0254P)^2 + 0.1986P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.37$ e Å⁻³

$\Delta\rho_{\min} = -0.35$ e Å⁻³

Extinction correction: SHELXL2014
(Sheldrick, 2015),

$F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.009 (2)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.0000	1.0000	1.0000	0.02186 (9)
O1	0.25606 (19)	1.00502 (12)	0.86692 (9)	0.0269 (2)
H1W	0.226 (4)	0.9463 (19)	0.8050 (8)	0.044 (6)*
H2W	0.303 (3)	1.0972 (10)	0.8515 (16)	0.041 (5)*
O2	0.2699 (2)	1.10732 (13)	1.11255 (11)	0.0345 (3)
H3W	0.307 (4)	1.2067 (5)	1.1282 (17)	0.048 (6)*
H4W	0.401 (2)	1.065 (3)	1.116 (2)	0.065 (7)*
O3	0.0981 (2)	0.78032 (12)	1.02590 (10)	0.0293 (2)
H5W	0.187 (3)	0.725 (2)	0.9825 (14)	0.051 (6)*
H6W	-0.008 (3)	0.7116 (19)	1.0423 (19)	0.059 (7)*
N1	0.5987 (2)	0.68143 (14)	0.16630 (10)	0.0246 (3)
N2	0.7365 (2)	0.58336 (15)	0.10516 (11)	0.0270 (3)
N3	0.6508 (2)	0.43523 (15)	0.10431 (11)	0.0296 (3)
N4	0.4544 (2)	0.43180 (14)	0.16488 (11)	0.0269 (3)
N5	-0.0918 (2)	0.58146 (17)	0.38266 (11)	0.0315 (3)
H5N	-0.180 (3)	0.5136 (17)	0.4067 (15)	0.042 (5)*
C1	0.4273 (2)	0.58463 (16)	0.20195 (12)	0.0212 (3)
C2	0.2376 (2)	0.64009 (17)	0.27458 (11)	0.0218 (3)
C3	0.2065 (3)	0.80168 (18)	0.30257 (13)	0.0284 (3)
H3A	0.3075	0.8778	0.2748	0.034*
C4	0.0250 (3)	0.8491 (2)	0.37189 (14)	0.0334 (4)
H4	0.0048	0.9570	0.3914	0.040*
C5	-0.1243 (3)	0.7363 (2)	0.41142 (14)	0.0345 (4)
H5	-0.2470	0.7668	0.4578	0.041*
C6	0.0827 (3)	0.53073 (19)	0.31686 (13)	0.0279 (3)
H6	0.0997	0.4219	0.2996	0.033*
N6	0.6295 (3)	0.22071 (16)	0.38657 (12)	0.0353 (3)
O4	0.5953 (2)	0.34960 (14)	0.44531 (11)	0.0442 (3)
O5	0.8117 (2)	0.21012 (15)	0.33009 (11)	0.0431 (3)
O6	0.4892 (4)	0.1062 (2)	0.38679 (19)	0.1001 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.02070 (14)	0.01556 (13)	0.02905 (15)	0.00093 (8)	0.00394 (9)	0.00145 (9)
O1	0.0295 (6)	0.0203 (5)	0.0304 (6)	-0.0002 (4)	0.0092 (4)	0.0015 (4)
O2	0.0292 (6)	0.0230 (6)	0.0485 (7)	-0.0009 (5)	-0.0073 (5)	-0.0031 (5)
O3	0.0283 (6)	0.0193 (5)	0.0419 (6)	0.0056 (4)	0.0149 (5)	0.0063 (5)
N1	0.0232 (6)	0.0217 (6)	0.0282 (6)	-0.0018 (5)	0.0064 (5)	0.0013 (5)

N2	0.0232 (6)	0.0257 (7)	0.0315 (7)	0.0011 (5)	0.0082 (5)	0.0013 (5)
N3	0.0282 (7)	0.0241 (6)	0.0360 (7)	0.0024 (5)	0.0095 (6)	0.0006 (5)
N4	0.0268 (7)	0.0199 (6)	0.0333 (7)	-0.0005 (5)	0.0086 (5)	0.0006 (5)
N5	0.0275 (7)	0.0374 (8)	0.0295 (7)	-0.0062 (6)	0.0086 (6)	0.0067 (6)
C1	0.0217 (7)	0.0197 (7)	0.0214 (7)	-0.0017 (5)	0.0015 (5)	0.0014 (5)
C2	0.0218 (7)	0.0228 (7)	0.0199 (7)	-0.0015 (5)	0.0018 (5)	0.0002 (5)
C3	0.0311 (8)	0.0239 (7)	0.0285 (8)	-0.0041 (6)	0.0076 (6)	-0.0004 (6)
C4	0.0373 (9)	0.0279 (8)	0.0332 (8)	0.0044 (7)	0.0079 (7)	-0.0049 (6)
C5	0.0283 (8)	0.0457 (10)	0.0284 (8)	0.0044 (7)	0.0092 (6)	-0.0015 (7)
C6	0.0279 (8)	0.0255 (8)	0.0297 (8)	-0.0030 (6)	0.0055 (6)	0.0033 (6)
N6	0.0398 (8)	0.0279 (7)	0.0346 (8)	-0.0094 (6)	0.0107 (6)	-0.0048 (6)
O4	0.0508 (8)	0.0301 (6)	0.0480 (7)	-0.0067 (5)	0.0237 (6)	-0.0080 (5)
O5	0.0428 (7)	0.0333 (6)	0.0502 (8)	-0.0024 (5)	0.0194 (6)	-0.0063 (6)
O6	0.0959 (14)	0.0607 (11)	0.1238 (16)	-0.0531 (10)	0.0672 (13)	-0.0446 (11)

Geometric parameters (Å, °)

Zn1—O3	2.0353 (11)	N4—C1	1.3345 (19)
Zn1—O3 ⁱ	2.0354 (11)	N5—C6	1.339 (2)
Zn1—O2	2.1011 (12)	N5—C5	1.339 (2)
Zn1—O2 ⁱ	2.1011 (12)	N5—H5N	0.8201 (11)
Zn1—O1 ⁱ	2.1841 (11)	C1—C2	1.463 (2)
Zn1—O1	2.1841 (11)	C2—C6	1.381 (2)
O1—H1W	0.8500	C2—C3	1.391 (2)
O1—H2W	0.8499	C3—C4	1.385 (2)
O2—H3W	0.8499	C3—H3A	0.9300
O2—H4W	0.8499	C4—C5	1.367 (2)
O3—H5W	0.8499	C4—H4	0.9300
O3—H6W	0.8501	C5—H5	0.9300
N1—C1	1.3384 (18)	C6—H6	0.9300
N1—N2	1.3405 (18)	N6—O6	1.210 (2)
N2—N3	1.3113 (18)	N6—O5	1.2485 (18)
N3—N4	1.3421 (18)	N6—O4	1.2525 (18)
O3—Zn1—O3 ⁱ	180.0	N2—N3—N4	109.65 (12)
O3—Zn1—O2	90.01 (5)	C1—N4—N3	104.64 (12)
O3 ⁱ —Zn1—O2	89.99 (5)	C6—N5—C5	122.90 (14)
O3—Zn1—O2 ⁱ	89.99 (5)	C6—N5—H5N	117.6 (14)
O3 ⁱ —Zn1—O2 ⁱ	90.01 (5)	C5—N5—H5N	119.5 (14)
O2—Zn1—O2 ⁱ	180.0	N4—C1—N1	111.53 (13)
O3—Zn1—O1 ⁱ	90.53 (4)	N4—C1—C2	124.52 (13)
O3 ⁱ —Zn1—O1 ⁱ	89.47 (4)	N1—C1—C2	123.93 (13)
O2—Zn1—O1 ⁱ	92.10 (5)	C6—C2—C3	118.26 (14)
O2 ⁱ —Zn1—O1 ⁱ	87.90 (5)	C6—C2—C1	119.94 (13)
O3—Zn1—O1	89.47 (4)	C3—C2—C1	121.80 (13)
O3 ⁱ —Zn1—O1	90.53 (4)	C4—C3—C2	119.93 (14)
O2—Zn1—O1	87.90 (5)	C4—C3—H3A	120.0
O2 ⁱ —Zn1—O1	92.10 (5)	C2—C3—H3A	120.0

O1 ⁱ —Zn1—O1	180.0	C5—C4—C3	119.67 (15)
Zn1—O1—H1W	118.9 (14)	C5—C4—H4	120.2
Zn1—O1—H2W	115.7 (13)	C3—C4—H4	120.2
H1W—O1—H2W	107.0 (18)	N5—C5—C4	119.30 (15)
Zn1—O2—H3W	126.7 (15)	N5—C5—H5	120.3
Zn1—O2—H4W	120.1 (17)	C4—C5—H5	120.3
H3W—O2—H4W	104 (2)	N5—C6—C2	119.93 (15)
Zn1—O3—H5W	123.7 (14)	N5—C6—H6	120.0
Zn1—O3—H6W	118.5 (15)	C2—C6—H6	120.0
H5W—O3—H6W	103 (2)	O6—N6—O5	120.33 (15)
C1—N1—N2	104.70 (12)	O6—N6—O4	120.02 (15)
N3—N2—N1	109.48 (11)	O5—N6—O4	119.63 (13)
C1—N1—N2—N3	0.11 (16)	N1—C1—C2—C3	7.1 (2)
N1—N2—N3—N4	-0.01 (17)	C6—C2—C3—C4	0.5 (2)
N2—N3—N4—C1	-0.09 (17)	C1—C2—C3—C4	-179.80 (14)
N3—N4—C1—N1	0.17 (17)	C2—C3—C4—C5	-0.7 (3)
N3—N4—C1—C2	-178.53 (13)	C6—N5—C5—C4	0.5 (3)
N2—N1—C1—N4	-0.18 (16)	C3—C4—C5—N5	0.2 (3)
N2—N1—C1—C2	178.53 (13)	C5—N5—C6—C2	-0.7 (2)
N4—C1—C2—C6	5.4 (2)	C3—C2—C6—N5	0.2 (2)
N1—C1—C2—C6	-173.17 (14)	C1—C2—C6—N5	-179.53 (13)
N4—C1—C2—C3	-174.32 (15)		

Symmetry code: (i) $-x, -y+2, -z+2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1W \cdots O5 ⁱⁱ	0.85	1.96	2.8067 (17)	172
O1—H2W \cdots N1 ⁱⁱⁱ	0.85	1.96	2.8029 (17)	173
O2—H3W \cdots N4 ^{iv}	0.85	2.02	2.8589 (17)	170
O2—H4W \cdots O1 ^v	0.85	2.08	2.9228 (17)	171
O3—H5W \cdots N3 ⁱⁱ	0.85	1.91	2.7446 (17)	168
O3—H6W \cdots N1 ^{vi}	0.85	2.72	3.4294 (17)	142
O3—H6W \cdots N2 ^{vi}	0.85	1.97	2.8076 (17)	169
N5—H5N \cdots N6 ^{vii}	0.82	2.61	3.344 (2)	149
N5—H5N \cdots O4 ^{vii}	0.82	1.92	2.7384 (18)	173
N5—H5N \cdots O5 ^{vii}	0.82	2.62	3.1347 (19)	123
C4—H4 \cdots O5 ^{viii}	0.93	2.65	3.452 (2)	145
C5—H5 \cdots O4 ^{ix}	0.93	2.52	3.292 (2)	141
C5—H5 \cdots O6 ^{ix}	0.93	2.52	3.422 (3)	165
C6—H6 \cdots O5 ^{vii}	0.93	2.41	3.047 (2)	126

Symmetry codes: (ii) $-x+1, -y+1, -z+1$; (iii) $-x+1, -y+2, -z+1$; (iv) $x, y+1, z+1$; (v) $-x+1, -y+2, -z+2$; (vi) $x-1, y, z+1$; (vii) $x-1, y, z$; (viii) $x-1, y+1, z$; (ix) $-x, -y+1, -z+1$.